

Age Does Matter: A Pilot Comparison of Placenta-Derived Stromal Cells for in utero Repair of Myelomeningocele Using a Lamb Model.

Journal: Fetal Diagn Ther

Publication Year: 2015

Authors: Erin G Brown, Benjamin A Keller, Lee Lankford, Christopher D Pivetti, Shinjiro Hirose, Diana L Farmer, Aijun Wang

PubMed link: 26159889

Funding Grants: UC Davis Stem Cell Training Program

Public Summary:

INTRODUCTION: Fetal amniotic membranes (FM) have been shown to preserve spinal cord histology in the fetal sheep model of myelomeningocele (MMC). This study compares the effectiveness of placenta-derived mesenchymal stromal cells (PMSCs) from early-gestation versus term-gestation placenta to augment FM repair to improve distal motor function in a sheep model. **METHODS:** Fetal lambs (n = 4) underwent surgical MMC creation followed by repair with FM patch with term-gestation PMSCs (n = 1), FM with early-gestation PMSCs (n = 1), FM only (n = 1), and skin closure only (n = 1). Histopathology and motor assessment was performed. **RESULTS:** Histopathologic analysis demonstrated increased preservation of spinal cord architecture and large neurons in the lamb repaired with early-gestation cells compared to all others. Lambs repaired with skin closure only, FM alone, and term-gestation PMSCs exhibited extremely limited distal motor function; the lamb repaired with early-gestation PMSCs was capable of normal ambulation. **DISCUSSION:** This pilot study is the first in vivo comparison of different gestational-age placenta-derived stromal cells for repair in the fetal sheep MMC model. The preservation of large neurons and markedly improved motor function in the lamb repaired with early-gestation cells suggest that early-gestation placental stromal cells may exhibit unique properties that augment in utero MMC repair to improve paralysis. © 2015 S. Karger AG, Basel.

Scientific Abstract:

INTRODUCTION: Fetal amniotic membranes (FM) have been shown to preserve spinal cord histology in the fetal sheep model of myelomeningocele (MMC). This study compares the effectiveness of placenta-derived mesenchymal stromal cells (PMSCs) from early-gestation versus term-gestation placenta to augment FM repair to improve distal motor function in a sheep model. **METHODS:** Fetal lambs (n = 4) underwent surgical MMC creation followed by repair with FM patch with term-gestation PMSCs (n = 1), FM with early-gestation PMSCs (n = 1), FM only (n = 1), and skin closure only (n = 1). Histopathology and motor assessment was performed. **RESULTS:** Histopathologic analysis demonstrated increased preservation of spinal cord architecture and large neurons in the lamb repaired with early-gestation cells compared to all others. Lambs repaired with skin closure only, FM alone, and term-gestation PMSCs exhibited extremely limited distal motor function; the lamb repaired with early-gestation PMSCs was capable of normal ambulation. **DISCUSSION:** This pilot study is the first in vivo comparison of different gestational-age placenta-derived stromal cells for repair in the fetal sheep MMC model. The preservation of large neurons and markedly improved motor function in the lamb repaired with early-gestation cells suggest that early-gestation placental stromal cells may exhibit unique properties that augment in utero MMC repair to improve paralysis. (c) 2015 S. Karger AG, Basel.

Source URL: <http://www.cirm.ca.gov/about-cirm/publications/age-does-matter-pilot-comparison-placenta-derived-stromal-cells-utero-repair>